

Datasheet for ABIN7303202

anti-CLIP3 antibody**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CLIP3
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CLIP3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (IHC), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CLIP3.
Specificity:	Recognizes endogenous levels of CLIP3 protein.
Characteristics:	Rabbit polyclonal antibody to CLIP3
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	CLIP3
Alternative Name:	CLIP3 (CLIP3 Products)
Background:	CLIPR59, CAP-Gly domain-containing linker protein 3, Cytoplasmic linker protein 170-related 59 kDa protein, CLIP-170-related 59 kDa protein, CLIPR-59

Target Details

Gene ID:	25999, 76686
UniProt:	Q96DZ5 , B9EHT4


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in PBS, pH 7.3, 0.2 % BSA, and 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of CLIP3 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

Image 2. Immunohistochemical analysis of CLIP3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated wi



Western Blotting

Image 3. Western blot analysis of CLIP3 expression in human brain (A), A549 (B) whole cell lysates.