

Datasheet for ABIN7303456

anti-TNS3 antibody**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	TNS3
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNS3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Tensin 3.
Specificity:	Recognizes endogenous levels of Tensin 3 protein.
Characteristics:	Rabbit polyclonal antibody to Tensin 3
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	TNS3
Alternative Name:	Tensin 3 (TNS3 Products)
Background:	TEM6, TENS1, TPP, Tensin-3, Tensin-like SH2 domain-containing protein 1, Tumor endothelial marker 6

Target Details

Gene ID: 64759

UniProt: [Q68CZ2](#)

Application Details

Application Notes: WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:100)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of Tensin 3 staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with t



Western Blotting

Image 2. Western blot analysis of Tensin 3 expression in HeLa (A), DLD (B), H9C2 (C) whole cell lysates.

Immunofluorescence

Image 3. Immunofluorescent analysis of Tensin 3 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibo