

Datasheet for ABIN7303590

**anti-EID1 antibody****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	EID1
Reactivity:	Human, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EID1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

## Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human EID1.
Specificity:	Recognizes endogenous levels of EID1 protein.
Characteristics:	Rabbit polyclonal antibody to EID1
Purification:	The antibody was purified by affinity chromatography.

## Target Details

Target:	EID1
Alternative Name:	EID1 ( <a href="#">EID1 Products</a> )
Background:	C15orf3, CRI1, RBP21, EP300-interacting inhibitor of differentiation 1, 21 kDa pRb-associated protein, CREBBP/EP300 inhibitory protein 1, E1A-like inhibitor of differentiation 1, EID-1

## Target Details

Gene ID: 23741

UniProt: [Q9Y6B2](#)

## Application Details

Application Notes: WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:100)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

## Images




### Immunofluorescence

**Image 1.** Immunofluorescent analysis of EID1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i



Immunohistochemistry

**Image 2.** Immunohistochemical analysis of EID1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the an



Western Blotting

**Image 3.** Western blot analysis of EID1 expression in PC3 (A), HeLa (B), HCT116 (C) whole cell lysates.