

Datasheet for ABIN7304384

**anti-NPY5R antibody****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	NPY5R
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NPY5R antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

## Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NPY5R.
Specificity:	Recognizes endogenous levels of NPY5R protein.
Characteristics:	Rabbit polyclonal antibody to NPY5R
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	NPY5R
Alternative Name:	NPY5R ( <a href="#">NPY5R Products</a> )
Background:	NPYR5, Neuropeptide Y receptor type 5, NPY5-R, NPY-Y5 receptor, NPY5-R, Y5 receptor

## Target Details

Gene ID:	4889
UniProt:	<a href="#">Q15761</a>
Pathways:	<a href="#">Feeding Behaviour</a>


## Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

## Images




### Immunofluorescence

**Image 1.** Immunofluorescent analysis of NPY5R staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

**Image 2.** Immunohistochemical analysis of NPY5R staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the a



Western Blotting

**Image 3.** Western blot analysis of NPY5R expression in Jurkat (A), HepG2 (B) whole cell lysates.