# antibodies .- online.com







# anti-HMGCR antibody

**Images** 



#### Overview

Quantity:	100 μL	
Target:	HMGCR	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HMGCR antibody is un-conjugated	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)	

# Product Details

Immunogen:	Recombinant full length protein of human HMGCR
Specificity:	Recognizes endogenous levels of HMGCR protein.
Characteristics:	Rabbit polyclonal antibody to HMGCR
Purification:	The antibody was purified by immunogen affinity chromatography.

# **Target Details**

Target:	HMGCR
Alternative Name:	HMGCR (HMGCR Products)
Background:	3-hydroxy-3-methylglutaryl-coenzyme A reductase, HMG-CoA reductase
Gene ID:	3156, 15357, 25675
UniProt:	P04035, Q01237, P51639

#### **Target Details**

Pathways:
-----------

AMPK Signaling, Negative Regulation of Hormone Secretion, Regulation of Lipid Metabolism by PPARalpha

#### **Application Details**


Application Notes:	WB (1:500 - 1:2000), IF/IC (1:50 - 1:200)
--------------------	---

Restrictions: For Research Use only

# Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



#### **Immunofluorescence**

Image 1. Immunofluorescent analysis of HMGCR staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



#### **Western Blotting**

Image 2. Western blot analysis of HMGCR expression in HepG2 (A), Hela (B), mouse spleen (C), rat liver (D) whole cell lysates.