antibodies -online.com





anti-PSMB9 antibody

3 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	PSMB9
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMB9 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	Recombinant full length protein of human PSMB9
Specificity:	Recognizes endogenous levels of PSMB9 protein.
Characteristics:	Rabbit polyclonal antibody to PSMB9
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	PSMB9
Alternative Name:	PSMB9 (PSMB9 Products)
Background:	LMP2, PSMB6i, RING12, Proteasome subunit beta type-9, Low molecular mass protein 2,
	Macropain chain 7, Multicatalytic endopeptidase complex chain 7, Proteasome chain 7, Proteasome subunit beta-1i, Really interesting new gene 12 protein

Target Details

Gene ID:	5698, 16912, 24967
UniProt:	P28065, P28076, P28077
Pathways:	Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA


Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:10 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months


Images



Immunofluorescence

Image 1. Immunofluorescent analysis of PSMB9 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody





Immunohistochemistry

Image 2. Immunohistochemical analysis of PSMB9 staining in mouse kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the

Western Blotting

Image 3. Western blot analysis of PSMB9 expression in HepG2 (A), THP1 (B), mouse liver (C) whole cell lysates.