antibodies .- online.com







anti-NPC2 antibody

Images



\sim					
()	VE	۲۱	/1	\triangle	Λ

Gene ID:

Quantity:	100 μL
Target:	NPC2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NPC2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)
Product Details	
Immunogen:	Recombinant full length protein of human NPC2
Specificity:	Recognizes endogenous levels of NPC2 protein.
Characteristics:	Rabbit polyclonal antibody to NPC2
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	NPC2
Alternative Name:	NPC2 (NPC2 Products)
Background:	HE1, Epididymal secretory protein E1, Human epididymis-specific protein 1, He1, Niemann-Pick disease type C2 protein

10577, 67963

Target Details

UniProt:	P61916, Q9Z0J0
Pathways:	SARS-CoV-2 Protein Interactome


Application Details

Application Notes:	WB (1:500 - 1:2000), IF/IC (1:50 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence

Image 1. Immunofluorescent analysis of NPC2 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i



Western Blotting

Image 2. Western blot analysis of NPC2 expression in HEK293T (A), mouse lung (B), mouse liver (C) whole cell lysates.