

Datasheet for ABIN7306508

anti-DLAT antibody**2** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DLAT
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DLAT antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	Recombinant full length protein of human PDC-E2
Specificity:	Recognizes endogenous levels of PDC-E2 protein.
Characteristics:	Rabbit polyclonal antibody to PDC-E2
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	DLAT
Alternative Name:	PDC-E2 (DLAT Products)
Background:	DLTA, Dihydrolipoyllysine-residue acetyltransferase component of pyruvate dehydrogenase complex, mitochondrial, 70 kDa mitochondrial autoantigen of primary biliary cirrhosis, PBC, Dihydrolipoamide acetyltransferase component of pyruvate dehydrogenase complex, M2 antigen complex 70 kDa subunit, Pyruvate dehydrogenase complex component E2, PDC-E2,

Target Details

	PDCE2
Gene ID:	1737, 235339, 81654
UniProt:	P10515 , Q8BMF4 , P08461


Application Details

Application Notes:	WB (1:500 - 1:2000), IF/IC (1:10 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence

Image 1. Immunofluorescent analysis of PDC-E2 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Western Blotting

Image 2. Western blot analysis of PDC-E2 expression in HepG2 (A), Jurkat (B), mouse heart (C), rat brain (D), rat kidney (E) whole cell lysates.