# antibodies -online.com





# anti-MDH2 antibody

3 Images



Go to Product page

## Overview

Quantity:	100 μL
Target:	MDH2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MDH2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

#### **Product Details**

Immunogen:	Recombinant full length protein of human MDH2
Specificity:	Recognizes endogenous levels of MDH2 protein.
Characteristics:	Rabbit polyclonal antibody to MDH2
Purification:	The antibody was purified by immunogen affinity chromatography.

# **Target Details**

Target:	MDH2
Alternative Name:	MDH2 (MDH2 Products)
Background:	Malate dehydrogenase, mitochondrial
Gene ID:	4191, 17448, 81829

# **Target Details**

UniProt:	P40926	P08249	P04636
Orni TOL.	1 70720	1 00277	, 1 0 7 0 0 0


# **Application Details**

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months


## **Images**



#### **Immunofluorescence**

**Image 1.** Immunofluorescent analysis of MDH2 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i





## **Immunohistochemistry**

**Image 2.** Immunohistochemical analysis of MDH2 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with

## **Western Blotting**

Image 3. Western blot analysis of MDH2 expression in Jurkat (A), A549 (B), mouse liver (C), mouse heart (D) whole cell lysates.