

Datasheet for ABIN7306654

**anti-KIF2B antibody**

3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	KIF2B
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIF2B antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

## Product Details

Immunogen:	Recombinant full length protein of human KIF2B
Specificity:	Recognizes endogenous levels of KIF2B protein.
Characteristics:	Rabbit polyclonal antibody to KIF2B
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	KIF2B
Alternative Name:	KIF2B ( <a href="#">KIF2B Products</a> )
Background:	Kinesin-like protein KIF2B
Gene ID:	84643, 73470, 287624

## Target Details

UniProt: [Q8N4N8](#), [Q8C0N1](#), [Q5XI51](#)

Pathways: [Microtubule Dynamics](#)

## Application Details

Application Notes: WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

## Images




### Immunofluorescence

**Image 1.** Immunofluorescent analysis of KIF2B staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

**Image 2.** Immunohistochemical analysis of KIF2B staining in mouse lung formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the an



Western Blotting

**Image 3.** Western blot analysis of KIF2B expression in U937 (A), mouse lung (B), rat lung (C) whole cell lysates.