



[Go to Product page](#)

Datasheet for ABIN7306740

## anti-MPP2 antibody

### 3 Images

#### Overview

Quantity:	100 µL
Target:	MPP2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MPP2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

#### Product Details

Immunogen:	Recombinant full length protein of human MPP2
Specificity:	Recognizes endogenous levels of MPP2 protein.
Characteristics:	Rabbit polyclonal antibody to MPP2
Purification:	The antibody was purified by immunogen affinity chromatography.

#### Target Details

Target:	MPP2
Alternative Name:	MPP2 ( <a href="#">MPP2 Products</a> )
Background:	DLG2, MAGUK p55 subfamily member 2, Discs large homolog 2, Protein MPP2
Gene ID:	4355, 50997

## Target Details

UniProt: [Q14168](#), [Q9WV34](#)

## Application Details

Application Notes: WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Expiry Date: 12 months

## Images




### Western Blotting

**Image 1.** Western blot analysis of MPP2 expression in HepG2 (A), mouse brain (B), rat brain (C) whole cell lysates.



### Immunofluorescence

**Image 2.** Immunofluorescent analysis of MPP2 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i



### Immunohistochemistry

**Image 3.** Immunohistochemical analysis of MPP2 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with