

Datasheet for ABIN7306770

anti-FARP2 antibody**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FARP2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FARP2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	Recombinant full length protein of human FIR
Specificity:	Recognizes endogenous levels of FIR protein.
Characteristics:	Rabbit polyclonal antibody to FIR
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	FARP2
Alternative Name:	FIR (FARP2 Products)
Background:	FIR, ROBPI, SIAHBP1, Poly(U)-binding-splicing factor PUF60, 60 kDa poly(U)-binding-splicing factor, FUSE-binding protein-interacting repressor, FBP-interacting repressor, Ro-binding protein 1, RoBP1, Siah-binding protein 1, Siah-BP1

Target Details

Gene ID:	22827, 67959
UniProt:	Q9UHX1 , Q3UEB3 , Q9WV25

Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of FIR staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in



Immunohistochemistry

Image 2. Immunohistochemical analysis of FIR staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the an



Western Blotting

Image 3. Western blot analysis of FIR expression in A549 (A), mouse brain (B), mouse spleen (C), rat lung (D), rat liver (E) whole cell lysates.