

Datasheet for ABIN7306770

**anti-FARP2 antibody****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	FARP2
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FARP2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

## Product Details

Immunogen:	Recombinant full length protein of human FIR
Specificity:	Recognizes endogenous levels of FIR protein.
Characteristics:	Rabbit polyclonal antibody to FIR
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	FARP2
Alternative Name:	FIR ( <a href="#">FARP2 Products</a> )
Background:	FIR, ROBPI, SIAHBP1, Poly(U)-binding-splicing factor PUF60, 60 kDa poly(U)-binding-splicing factor, FUSE-binding protein-interacting repressor, FBP-interacting repressor, Ro-binding protein 1, RoBP1, Siah-binding protein 1, Siah-BP1

## Target Details

Gene ID:	22827, 67959
UniProt:	<a href="#">Q9UHX1</a> , <a href="#">Q3UEB3</a> , <a href="#">Q9WV25</a>


## Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

## Images




### Immunofluorescence

**Image 1.** Immunofluorescent analysis of FIR staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in



Immunohistochemistry

**Image 2.** Immunohistochemical analysis of FIR staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the an



Western Blotting

**Image 3.** Western blot analysis of FIR expression in A549 (A), mouse brain (B), mouse spleen (C), rat lung (D), rat liver (E) whole cell lysates.