# antibodies .- online.com





anti-VSNL1 antibody



**Images** 



#### Go to Product page

$\sim$				٠			
	۱۱/	Δ	rv	I		1/	1
$\cup$	v	C	ΙV	ı	$\overline{}$	٧	V

Quantity:	100 μL
Target:	VSNL1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VSNL1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

#### **Product Details**

Immunogen:	Recombinant full length protein of human VILIP-1
Specificity:	Recognizes endogenous levels of VILIP-1 protein.
Characteristics:	Rabbit polyclonal antibody to VILIP-1
Purification:	The antibody was purified by immunogen affinity chromatography.

### **Target Details**

Target:	VSNL1
Alternative Name:	VILIP-1 (VSNL1 Products)
Background:	VISL1, Visinin-like protein 1, VILIP, VLP-1, Hippocalcin-like protein 3, HLP3
Gene ID:	7447, 26950

#### **Target Details**

UniProt:	P62760, P62761, P62762


## **Application Details**

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:100)
Restrictions:	For Research Use only

#### Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**




#### **Immunofluorescence**

**Image 1.** Immunofluorescent analysis of VILIP-1 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibod



#### **Immunohistochemistry**

**Image 2.** Immunohistochemical analysis of VILIP-1 staining in mouse brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the



#### **Western Blotting**

Image 3. Western blot analysis of VILIP-1 expression in IMR32 (A) whole cell lysates.