antibodies -online.com





anti-TRIP10 antibody





Go to Product page

\sim	
()\/\	view
	V I C V V

Quantity:100 μLTarget:TRIP10Reactivity:Human, MouseHost:RabbitClonality:PolyclonalConjugate:This TRIP10 antibody is un-conjugatedApplication:Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)	0.0	
Reactivity: Human, Mouse Host: Rabbit Clonality: Polyclonal Conjugate: This TRIP10 antibody is un-conjugated	Quantity:	100 μL
Host: Rabbit Clonality: Polyclonal Conjugate: This TRIP10 antibody is un-conjugated	Target:	TRIP10
Clonality: Polyclonal Conjugate: This TRIP10 antibody is un-conjugated	Reactivity:	Human, Mouse
Conjugate: This TRIP10 antibody is un-conjugated	Host:	Rabbit
	Clonality:	Polyclonal
Application: Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)	Conjugate:	This TRIP10 antibody is un-conjugated
	Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)
Product Dataile	Droduct Dotaile	

Product Details

Immunogen:	Recombinant full length protein of human CIP4
Specificity:	Recognizes endogenous levels of CIP4 protein.
Characteristics:	Rabbit polyclonal antibody to CIP4
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	TRIP10
Alternative Name:	CIP4 (TRIP10 Products)
Background:	CIP4, STOT, STP, Cdc42-interacting protein 4, Protein Felic, Salt tolerant protein, hSTP, Thyroid receptor-interacting protein 10, TR-interacting protein 10, TRIP-10
Gene ID:	9322, 106628

Target Details

UniProt: Q15642, Q8CJ53


Application Details

Application Notes:	WB (1:500 - 1:2000), IF/IC (1:50 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence

Image 1. Immunofluorescent analysis of CIP4 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i



Western Blotting

Image 2. Western blot analysis of CIP4 expression in Hela (A), SKOV3 (B), mouse kidney (C), mouse heart (D) whole cell lysates.