antibodies -online.com





anti-Transglutaminase 5 antibody

3 Images



Go to Product page

Overview

Quantity:	100 μL
Target:	Transglutaminase 5 (TGM5)
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Transglutaminase 5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	Recombinant full length protein of human TGase5
Specificity:	Recognizes endogenous levels of TGase5 protein.
Characteristics:	Rabbit polyclonal antibody to TGase5
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	Transglutaminase 5 (TGM5)
Alternative Name:	TGase5 (TGM5 Products)
Background:	TGMX, Protein-glutamine gamma-glutamyltransferase 5, Transglutaminase X, TG(X), TGX, TGase X, Transglutaminase-5, TGase-5

Target Details

Gene ID:	9333, 74176
UniProt:	O43548, Q9D7I9


Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence

Image 1. Immunofluorescent analysis of TGase5 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



80

60

kDa В 200 140 100

Immunohistochemistry

Image 2. Immunohistochemical analysis of TGase5 staining in rat kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the a

Western Blotting

Image 3. Western blot analysis of TGase5 expression in mouse liver (A), rat heart (B) whole cell lysates.