

Datasheet for ABIN7306928

anti-STAMPB antibody**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	STAMPB
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAMPB antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	Recombinant full length protein of human STAMPB
Specificity:	Recognizes endogenous levels of STAMPB protein.
Characteristics:	Rabbit polyclonal antibody to STAMPB
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	STAMPB
Alternative Name:	STAMPB (STAMPB Products)
Background:	AMSH, STAM-binding protein, Associated molecule with the SH3 domain of STAM, Endosome-associated ubiquitin isopeptidase

Target Details

Gene ID: 10617, 70527, 171565

UniProt: [O95630](#), [Q9CQ26](#), [Q8R424](#)

Application Details

Application Notes: WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:100)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of STAMBP staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

Image 2. Immunohistochemical analysis of STAMBP staining in rat spleen formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the a



Western Blotting

Image 3. Western blot analysis of STAMBP expression in HepG2 (A), Hela (B), mouse brain (C), rat lung (D) whole cell lysates.