

Datasheet for ABIN7306932

**anti-SUB1 antibody**

3 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	SUB1
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SUB1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

## Product Details

Immunogen:	Recombinant full length protein of human PC4
Specificity:	Recognizes endogenous levels of PC4 protein.
Characteristics:	Rabbit polyclonal antibody to PC4
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	SUB1
Alternative Name:	PC4 ( <a href="#">SUB1 Products</a> )
Background:	PC4, RPO2TC1, Activated RNA polymerase II transcriptional coactivator p15, Positive cofactor 4, PC4, SUB1 homolog, p14

## Target Details

Gene ID: 10923, 20024

UniProt: [P53999](#), [P11031](#)

## Application Details

Application Notes: WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:100)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

## Images




### Immunofluorescence

**Image 1.** Immunofluorescent analysis of PC4 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in



Immunohistochemistry

**Image 2.** Immunohistochemical analysis of PC4 staining in human stomach formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the a



Western Blotting

**Image 3.** Western blot analysis of PC4 expression in HL60 (A), Jurkat (B), mouse brain (C) whole cell lysates.