

Datasheet for ABIN7306990

anti-HMG20A antibody**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HMG20A
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMG20A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	Recombinant full length protein of human HMG-20A
Specificity:	Recognizes endogenous levels of HMG-20A protein.
Characteristics:	Rabbit polyclonal antibody to HMG-20A
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	HMG20A
Alternative Name:	HMG-20A (HMG20A Products)
Background:	HMGX1, HMGXB1, High mobility group protein 20A, HMG box-containing protein 20A, HMG domain-containing protein 1, HMG domain-containing protein HMGX1

Target Details

Gene ID:	10363, 66867
UniProt:	Q9NP66 , Q9DC33


Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images




Western Blotting

Image 1. Western blot analysis of HMG-20A expression in HEK293T (A), HeLa (B) whole cell lysates.



Immunofluorescence

Image 2. Immunofluorescent analysis of HMG-20A staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

Image 3. Immunohistochemical analysis of HMG-20A staining in mouse brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the