antibodies - online.com







anti-CCAR1 antibody





\sim						
	1//	Д	r۱	1	\triangle	٨

Quantity:	100 μL
Target:	CCAR1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CCAR1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Purpose:	Rabbit polyclonal antibody to CCAR1
Immunogen:	Recombinant full length protein of human CCAR1
Specificity:	Recognizes endogenous levels of CCAR1 protein.
Characteristics:	Rabbit polyclonal antibody to CCAR1
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	CCAR1
Alternative Name:	CCAR1 (CCAR1 Products)
Background:	CARP1, DIS, Cell division cycle and apoptosis regulator protein 1, Cell cycle and apoptosis
	regulatory protein 1, CARP-1, Death inducer with SAP domain

Target Details

Gene ID:	55749, 67500
UniProt:	Q8IX12, Q8CH18


Application Details

Application Notes:	WB (1:500 - 1:2000), IF/IC (1:10 - 1:100)	
Restrictions:	For Research Use only	

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence

Image 1. Immunofluorescent analysis of CCAR1 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Western Blotting

Image 2. Western blot analysis of CCAR1 expression in Jurkat (A), HepG2 (B), A549 (C), A431 (D) whole cell lysates.