



[Go to Product page](#)

Datasheet for ABIN7307660
anti-CIAPIN1 antibody

3 Images

Overview

Quantity:	100 µL
Target:	CIAPIN1
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CIAPIN1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Purpose:	Rabbit polyclonal antibody to Anamorsin
Immunogen:	Recombinant full length protein of human Anamorsin
Specificity:	Recognizes endogenous levels of Anamorsin protein.
Characteristics:	Rabbit polyclonal antibody to Anamorsin
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	CIAPIN1
Alternative Name:	Anamorsin (CIAPIN1 Products)
Background:	Anamorsin, Cytokine-induced apoptosis inhibitor 1, Fe-S cluster assembly protein DRE2

Target Details

	homolog
Gene ID:	57019, 109006
UniProt:	Q6F181 , Q8WTY4


Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of Anamorsin staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary anti



Immunohistochemistry

Image 2. Immunohistochemical analysis of Anamorsin staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated



Western Blotting

Image 3. Western blot analysis of Anamorsin expression in HepG2 (A), Hela (B), MCF7 (C), mouse testis (D) whole cell lysates.