



[Go to Product page](#)

Datasheet for ABIN7307738
anti-DLGAP5 antibody

3 Images

Overview

Quantity:	100 µL
Target:	DLGAP5
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DLGAP5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (IHC), Immunochromatography (IC)

Product Details

Purpose:	Rabbit polyclonal antibody to DLGAP5
Immunogen:	Recombinant full length protein of human DLGAP5
Specificity:	Recognizes endogenous levels of DLGAP5 protein.
Characteristics:	Rabbit polyclonal antibody to DLGAP5
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	DLGAP5
Alternative Name:	DLGAP5 (DLGAP5 Products)
Background:	DLG7, KIAA0008, Disks large-associated protein 5, DAP-5, Discs large homolog 7, Disks large-

Target Details

associated protein DLG7, Hepatoma up-regulated protein, HURP

Gene ID: 9787, 218977

UniProt: [Q15398](#), [Q8K4R9](#)

Pathways: [M Phase](#)

Application Details

Application Notes: WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of DLGAP5 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

Image 2. Immunohistochemical analysis of DLGAP5 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated w



Western Blotting

Image 3. Western blot analysis of DLGAP5 expression in mouse brain (A), mouse liver (B) whole cell lysates.