

Datasheet for ABIN7307740

anti-DNAJB6 antibody**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DNAJB6
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DNAJB6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Purpose:	Rabbit polyclonal antibody to DNAJB6
Immunogen:	Recombinant full length protein of human DNAJB6
Specificity:	Recognizes endogenous levels of DNAJB6 protein.
Characteristics:	Rabbit polyclonal antibody to DNAJB6
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	DNAJB6
Alternative Name:	DNAJB6 (DNAJB6 Products)
Background:	HSJ2, MRJ, MSJ1, DnaJ homolog subfamily B member 6, HHDJ1, Heat shock protein J2, HSJ-

Target Details

	2, MRJ, MSJ-1
Gene ID:	10049, 23950, 362293
UniProt:	O75190 , O54946 , Q6AYU3


Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of DNAJB6 staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the



Western Blotting

Image 2. Western blot analysis of DNAJB6 expression in MCF7 (A), Jurkat (B), HeLa (C), mouse testis (D), rat brain (E) whole cell lysates.



Immunofluorescence

Image 3. Immunofluorescent analysis of DNAJB6 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody