

Datasheet for ABIN7307868

**anti-HDGF antibody****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	HDGF
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HDGF antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Immunochromatography (IC)

## Product Details

Purpose:	Rabbit polyclonal antibody to HDGF
Immunogen:	Recombinant full length protein of human HDGF
Specificity:	Recognizes endogenous levels of HDGF protein.
Characteristics:	Rabbit polyclonal antibody to HDGF
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	HDGF
Alternative Name:	HDGF ( <a href="#">HDGF Products</a> )
Background:	HMG1L2, Hepatoma-derived growth factor, HDGF, High mobility group protein 1-like 2, HMG-

## Target Details

	1L2
Gene ID:	3068, 15191, 114499
UniProt:	<a href="#">P51858</a> , <a href="#">P51859</a> , <a href="#">Q8VHK7</a>
Pathways:	<a href="#">ER-Nucleus Signaling</a>

## Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200), IP (1:20 - 1:50)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

## Images




### Immunofluorescence

**Image 1.** Immunofluorescent analysis of HDGF staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody i



**Immunohistochemistry**

**Image 2.** Immunohistochemical analysis of HDGF staining in rat brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the anti



**Western Blotting**

**Image 3.** Western blot analysis of HDGF expression in Hela (A), A549 (B), MCF7 (C) whole cell lysates.