

Datasheet for ABIN7307932

anti-IL23R antibody

3 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IL23R
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL23R antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Purpose:	Rabbit polyclonal antibody to IL-23R
Immunogen:	Recombinant full length protein of human IL-23R
Specificity:	Recognizes endogenous levels of IL-23R protein.
Characteristics:	Rabbit polyclonal antibody to IL-23R
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	IL23R
Alternative Name:	IL-23R (IL23R Products)
Background:	Interleukin-23 receptor, IL-23 receptor, IL-23R

Target Details

Gene ID:	149233, 209590
UniProt:	Q5VWK5 , Q8K4B4
Pathways:	Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Activated T Cell Proliferation


Application Details

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of IL-23R staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

Image 2. Immunohistochemical analysis of IL-23R staining in human spleen formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the



Western Blotting

Image 3. Western blot analysis of IL-23R expression in K562 (A), HepG2 (B), mouse intestine (C), mouse spleen (D) whole cell lysates.