# antibodies -online.com









#### Overview

Quantity:	100 μL
Target:	INTS6
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This INTS6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

#### **Product Details**

Purpose:	Rabbit polyclonal antibody to INTS6
Immunogen:	Recombinant full length protein of human INTS6
Specificity:	Recognizes endogenous levels of INTS6 protein.
Characteristics:	Rabbit polyclonal antibody to INTS6
Purification:	The antibody was purified by immunogen affinity chromatography.

## **Target Details**

Target:	INTS6
Alternative Name:	INTS6 (INTS6 Products)
Background:	DBI1, DDX26, DDX26A, Integrator complex subunit 6, Int6, DBI-1, Protein DDX26, Protein deleted

## **Target Details**

	in cancer 1, DICE1
Gene ID:	26512, 18130
UniProt:	Q9UL03, Q6PCM2


# **Application Details**

Application Notes:	WB (1:500 - 1:2000), IH (1:50 - 1:200), IF/IC (1:50 - 1:200)
Restrictions:	For Research Use only

# Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

### **Images**




## **Immunohistochemistry**

**Image 1.** Immunohistochemical analysis of INTS6 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated wi



#### **Western Blotting**

Image 2. Western blot analysis of INTS6 expression in HL60 (A), SW620 (B), Hela (C) whole cell lysates.



#### **Immunofluorescence**

**Image 3.** Immunofluorescent analysis of INTS6 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody