# antibodies -online.com





# anti-MYH1 antibody

2 Images



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	MYH1
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MYH1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)

## **Product Details**

Purpose:	Rabbit polyclonal antibody to Myosin 1
Immunogen:	Recombinant full length protein of human Myosin 1
Specificity:	Recognizes endogenous levels of Myosin 1 protein.
Characteristics:	Rabbit polyclonal antibody to Myosin 1
Purification:	The antibody was purified by immunogen affinity chromatography.

# Target Details

Target:	MYH1
Alternative Name:	Myosin 1 (MYH1 Products)
Background:	Myosin-1, Myosin heavy chain 1, Myosin heavy chain 2x, MyHC-2x, Myosin heavy chain IIx/d, MyHC-IIx/d, Myosin heavy chain, skeletal muscle, adult 1

## **Target Details**

Gene ID:	4619, 17879
UniProt:	P12882, Q5SX40


## **Application Details**

Application Notes:	WB (1:500 - 1:2000), IF/IC (1:50 - 1:100)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



#### **Immunofluorescence**

**Image 1.** Immunofluorescent analysis of Myosin 1 staining in U2OS cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibo



#### **Western Blotting**

Image 2. Western blot analysis of Myosin 1 expression in mouse heart (A), rat skeletal muscle (B), rat heart (C) whole cell lysates.