# antibodies .- online.com





## anti-POLR1D antibody

2 Images



Go to Product page

$\sim$							
	1//	$\Box$	$r \setminus$	/ [	$\bigcirc$	1	٨,

Background:

Quantity:	100 μL
Target:	POLR1D
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This POLR1D antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunochromatography (IC)
Product Details	
Purpose:	Rabbit polyclonal antibody to POLR1D
Purpose: Immunogen:	Rabbit polyclonal antibody to POLR1D  Recombinant full length protein of human POLR1D
·	
Immunogen:	Recombinant full length protein of human POLR1D
Immunogen: Specificity:	Recombinant full length protein of human POLR1D  Recognizes endogenous levels of POLR1D protein.
Immunogen: Specificity: Characteristics:	Recombinant full length protein of human POLR1D  Recognizes endogenous levels of POLR1D protein.  Rabbit polyclonal antibody to POLR1D
Immunogen:  Specificity:  Characteristics:  Purification:	Recombinant full length protein of human POLR1D  Recognizes endogenous levels of POLR1D protein.  Rabbit polyclonal antibody to POLR1D

DNA-directed RNA polymerases I and III subunit RPAC2, RNA polymerases I and III subunit AC2,

AC19, DNA-directed RNA polymerase I subunit D, RNA polymerase I 16 kDa subunit, RPA16,

### **Target Details**

	RPC16, hRPA19
Gene ID:	51082
UniProt:	Q9Y2S0


### **Application Details**

Application Notes:	WB (1:500 - 1:2000), IF/IC (1:50 - 1:100)	
Restrictions:	For Research Use only	

### Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



#### **Immunofluorescence**

**Image 1.** Immunofluorescent analysis of POLR1D staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



#### **Western Blotting**

**Image 2.** Western blot analysis of POLR1D expression in MCF7 (A), A549 (B) whole cell lysates.