Datasheet for ABIN731201 anti-SLC8A1 antibody (AA 801-900) (Cy3)

-online.com antibodies



Overview

Quantity:	100 μL	
Target:	SLC8A1	
Binding Specificity:	AA 801-900	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SLC8A1 antibody is conjugated to Cy3	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NCX1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Chicken,Guinea Pig
Purification:	Purified by Protein A.
Target Details	
Target:	SLC8A1
Alternative Name:	NCX1/SLC8A1 (SLC8A1 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN731201 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Background:	Synonyms: Na+/Ca2+exchanger 1, CNC, DKFZp779F0871, MGC119581 , FLJ37694, FLJ43417,		
	Na+/Ca2+ exchange protein 1, Na+/Ca2+ exchanger, NCX 1, NCX, SLC8A1, SLC8A1 protein ,		
	Sodium Calcium Exchanger, Sodium/calcium exchanger 1, Solute carrier family 8 member 1.		
	Background: In cardiac myocytes, Ca(2+) concentrations alternate between high levels during		
	contraction and low levels during relaxation. The increase in Ca(2+) concentration during		
	contraction is primarily due to release of Ca(2+) from intracellular stores. However, some		
	Ca(2+) also enters the cell through the sarcolemma(plasma membrane). During relaxation,		
	Ca(2+) is sequestered within the intracellular stores. To prevent overloading of intracellular		
	stores, the Ca(2+) that entered across the sarcolemma must be extruded from the cell. The		
	Na(+)-Ca(2+) exchanger is the primary mechanism by which the Ca(2+) is extruded from the		
	cell during relaxation. In the heart, the exchanger may play a key role in digitalis action. The		
	exchanger is the dominant mechanism in returning the cardiac myocyte to its resting state		
	following excitation.[supplied by OMIM].		
Gene ID:	6546		
Pathways:	Myometrial Relaxation and Contraction		

Application Details

Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN731201 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

lond	lina
land	
10110	

Storage Comment:

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN731201 | 03/07/2024 | Copyright antibodies-online. All rights reserved.