# antibodies -online.com





## anti-SLC8A1 antibody (AA 801-900) (Cy5)



$\sim$	4.4	D		page
1-()	17	$\mathbf{P}_{\mathbf{r}}$	וי או וו	nana
$\cup$	w	1 100	IUCL	Dauc

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	SLC8A1
Binding Specificity:	AA 801-900
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC8A1 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human NCX1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Chicken,Guinea Pig
Purification:	Purified by Protein A.

### Target Details

Target:	SLC8A1
Alternative Name:	NCX1/SLC8A1 (SLC8A1 Products)

### Target Details

Background:	Synonyms: Na+/Ca2+exchanger 1, CNC, DKFZp779F0871, MGC119581 , FLJ37694, FLJ43417,
	Na+/Ca2+ exchange protein 1, Na+/Ca2+ exchanger, NCX 1, NCX, SLC8A1, SLC8A1 protein ,
	Sodium Calcium Exchanger, Sodium/calcium exchanger 1, Solute carrier family 8 member 1.
	Background: In cardiac myocytes, Ca(2+) concentrations alternate between high levels during
	contraction and low levels during relaxation. The increase in Ca(2+) concentration during
	contraction is primarily due to release of Ca(2+) from intracellular stores. However, some
	Ca(2+) also enters the cell through the sarcolemma(plasma membrane). During relaxation,
	Ca(2+) is sequestered within the intracellular stores. To prevent overloading of intracellular
	stores, the Ca(2+) that entered across the sarcolemma must be extruded from the cell. The
	Na(+)-Ca(2+) exchanger is the primary mechanism by which the Ca(2+) is extruded from the
	cell during relaxation. In the heart, the exchanger may play a key role in digitalis action. The
	exchanger is the dominant mechanism in returning the cardiac myocyte to its resting state
	following excitation.[supplied by OMIM].
Gene ID:	6546
Pathways:	Myometrial Relaxation and Contraction
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

## Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months