

Datasheet for ABIN731207

anti-SLC8A1 antibody (AA 801-900) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	SLC8A1
Binding Specificity:	AA 801-900
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC8A1 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NCX1
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Horse,Chicken,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	SLC8A1
Alternative Name:	NCX1/SLC8A1 (SLC8A1 Products)

Target Details

Background:	<p>Synonyms: Na⁺/Ca²⁺exchanger 1, CNC, DKFZp779F0871, MGC119581 , FLJ37694, FLJ43417, Na⁺/Ca²⁺ exchange protein 1, Na⁺/Ca²⁺ exchanger, NCX 1, NCX, SLC8A1, SLC8A1 protein , Sodium Calcium Exchanger, Sodium/calcium exchanger 1, Solute carrier family 8 member 1.</p> <p>Background: In cardiac myocytes, Ca(2+) concentrations alternate between high levels during contraction and low levels during relaxation. The increase in Ca(2+) concentration during contraction is primarily due to release of Ca(2+) from intracellular stores. However, some Ca(2+) also enters the cell through the sarcolemma(plasma membrane). During relaxation, Ca(2+) is sequestered within the intracellular stores. To prevent overloading of intracellular stores, the Ca(2+) that entered across the sarcolemma must be extruded from the cell. The Na(+)-Ca(2+) exchanger is the primary mechanism by which the Ca(2+) is extruded from the cell during relaxation. In the heart, the exchanger may play a key role in digitalis action. The exchanger is the dominant mechanism in returning the cardiac myocyte to its resting state following excitation.[supplied by OMIM].</p>
-------------	---

Gene ID:	6546
Pathways:	Myometrial Relaxation and Contraction

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months