antibodies -online.com





Glucagon Protein (GCG) (AA 21-180) (His tag)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	50 µg
Target:	Glucagon (GCG)
Protein Characteristics:	AA 21-180
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Glucagon protein is labelled with His tag.
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Mass Spectrometry (MS)
Product Details	
Purpose:	Recombinant Human GCG Protein is produced by our Mammalian expression system and the
	target gene encoding Arg21-Lys180 is expressed with a 6His tag at the C-terminus.
Sequence:	RSLQDTEEKS RSFSASQADP LSDPDQMNED KRHSQGTFTS DYSKYLDSRR AQDFVQWLMN
	TKRNRNNIAK RHDEFERHAE GTFTSDVSSY LEGQAAKEFI AWLVKGRGRR DFPEEVAIVE
	ELGRRHADGS FSDEMNTILD NLAARDFINW LIQTKITDRK
Purity:	Greater than 95 % as determined by reducing SDS-PAGE.
Sterility:	0.2 μm filtered
Endotoxin Level:	Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test.
Target Details	
Target:	Glucagon (GCG)

Target Details

Alternative Name:	GCG (GCG Products)	
Gene ID:	2641	
UniProt:	P01275	
Pathways:	Positive Regulation of Peptide Hormone Secretion, Peptide Hormone Metabolism, cAMP Metabolic Process, Regulation of Carbohydrate Metabolic Process, Feeding Behaviour, Negative Regulation of intrinsic apoptotic Signaling	

For Research Use only

Application Details

Restrictions:

Handling	
Format:	Liquid
Buffer:	Supplied as a 0.2 μ M filtered solution of 20 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 50 % Glycerol, pH 8.0.
Handling Advice:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in 1X PBS. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.