

Datasheet for ABIN7313154 PDGFA Protein (AA 87-211)



Overview

Quantity:	50 µg
Target:	PDGFA
Protein Characteristics:	AA 87-211
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Application:	Western Blotting (WB), SDS-PAGE (SDS), ELISA, Mass Spectrometry (MS)
Product Details	
Purpose:	Recombinant Human PDGFA Protein is produced by our E. coli expression system and the
	target gene encoding Ser87-Thr211 is expressed.
Sequence:	SIEEAVPAVC KTRTVIYEIP RSQVDPTSAN FLIWPPCVEV KRCTGCCNTS SVKCQPSRVH
	HRSVKVAKVE YVRKKPKLKE VQVRLEEHLE CACATTSLNP DYREEDTGRP RESGKKRKRK RLKPT
Purity:	Greater than 95 % as determined by reducing SDS-PAGE.
Sterility:	0.2 µm filtered
Endotoxin Level:	Less than 0.1 ng/µg (1 IEU/µg) as determined by LAL test.
Target Details	
i ai yet Detalis	

Target:	PDGFA
Alternative Name:	PDGFA (PDGFA Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7313154 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Gene ID:	5154
UniProt:	P04085
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Smooth Muscle Cell Migration, Platelet-derived growth Factor Receptor Signaling
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	Lyophilized from a 0.2 μ M filtered solution of PBS, pH 7.4.
Handling Advice:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in 1X PBS. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.