



[Go to Product page](#)

Datasheet for ABIN7314182
DcR2 Protein (AA 56-211) (Fc Tag)

Overview

Quantity:	50 µg
Target:	DcR2 (TNFRSF10D)
Protein Characteristics:	AA 56-211
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This DcR2 protein is labelled with Fc Tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Mass Spectrometry (MS)

Product Details

Purpose:	Recombinant Human CD264 Protein is produced by mammalian expression system and the target gene encoding Ala56-His211 is expressed with a Fc tag at the C-terminus.
Sequence:	<p>ATIPRQDEVP QQT VAPQQQR RSLKEEECPA GSHRSEYTG A CNPCTEGVDY TIASNNLPSC</p> <p>LLCTVCKSGQ TNKSSCTTTR DTVCQCEKGS FQDKNSPEMC RTCRTGCPRG MVKVSNCTPR</p> <p>SDIKCKNESA ASSTGKTPAA EETVTTILGM LASYHIEGR IDPKSCDKTH TCPPCPAPEL</p> <p>LGGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV HNAKTKPREE</p> <p>QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKALPAPIEK TISKAKGQPR EPQVYTLPPS</p> <p>RDELTKNQVS LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF FLYSKLTVDK</p> <p>SRWQQGNVFS CSVMHEALHN HYTQKSLSL S PGK</p>
Purity:	Greater than 95 % as determined by reducing SDS-PAGE.
Sterility:	0.2 µm filtered

Target Details

Target:	DcR2 (TNFRSF10D)
Alternative Name:	TNFRSF10D (TNFRSF10D Products)
Gene ID:	8793
UniProt:	Q9UBN6
Pathways:	Apoptosis

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Lyophilized
Buffer:	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Handling Advice:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH ₂ O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.