.-online.com antibodies

## Datasheet for ABIN7314420 IL28A Protein (AA 26-200) (His tag)



Overview	
Quantity:	50 µg
Target:	IL28A
Protein Characteristics:	AA 26-200
Origin:	Human
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL28A protein is labelled with His tag.
Application:	SDS-PAGE (SDS), Western Blotting (WB), ELISA, Mass Spectrometry (MS)
Product Details	
Purpose:	Recombinant Human IL-28A Protein is produced by mammalian expression system and the
	target gene encoding Val26-Val200 is expressed with a 6His tag at the C-terminus.
Sequence:	VPVARLHGAL PDARGCHIAQ FKSLSPQELQ AFKRAKDALE ESLLLKDCRC HSRLFPRTWD
	LRQLQVRERP MALEAELALT LKVLEATADT DPALVDVLDQ PLHTLHHILS QFRACIQPQP
	TAGPRTRGRL HHWLYRLQEA PKKESPGCLE ASVTFNLFRL LTRDLNCVAS GDLCVHHHHH H
Purity:	Greater than 95 % as determined by reducing SDS-PAGE.
Sterility:	0.2 µm filtered
Target Details	
Target:	IL28A

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7314420 | 04/11/2024 | Copyright antibodies-online. All rights reserved.

IFNL2 (IL28A Products)

Alternative Name:

## Target Details

Gene ID:	282616
UniProt:	Q8IZJ0

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Handling Advice:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH20. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.