-online.com antibodies

## Datasheet for ABIN7314720 SLAMF8 Protein (AA 21-231) (His tag)



Overview	
Quantity:	50 µg
Target:	SLAMF8
Protein Characteristics:	AA 21-231
Origin:	Mouse
Source:	Human Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SLAMF8 protein is labelled with His tag.
Application:	ELISA, SDS-PAGE (SDS), Western Blotting (WB), Mass Spectrometry (MS)
Product Details	
Purpose:	Recombinant Mouse CD353 Protein is produced by mammalian expression system and the
	target gene encoding Val21-Asp231 is expressed with a 6His tag at the C-terminus.
Sequence:	VQVLSKVGDS ELLVAECPPG FQVREAIWRS LWPSEELLAT FFRGSLETLY HSRFLGRVQL
	YDNLSLELGP LKPGDSGNFS VLMVDTGGQT WTQTLYLKVY DAVPKPEVQV FTAAAEETQP
	LNTCQVFLSC WAPNISDITY SWRWEGTVDF NGEVRSHFSN GQVLSVSLGL GDKDVAFTCI
	ASNPVSWDMT TVTPWESCHH EAASGKASYK DHHHHHH
Purity:	Greater than 95 % as determined by reducing SDS-PAGE.
Sterility:	0.2 µm filtered
Target Details	

Target:

SLAMF8

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7314720 | 04/11/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	SLAMF8 (SLAMF8 Products)
Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Buffer:	Lyophilized from a 0.2 $\mu m$ filtered solution of PBS, $$ pH 7.4.
Handling Advice:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 µg/mL. Dissolve the lyophilized protein in ddH20. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	Lyophilized protein should be stored at -20°C, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at -20°C for 3 months.