antibodies .- online.com





SARS-CoV-2 Spike S1 Protein (BA.2.75 - Omicron) (His tag)



Of to Freduct page

()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	I١	νı	ㄷ	٧	۷

Quantity:	100 μg
Target:	SARS-CoV-2 Spike S1
Protein Characteristics:	BA.2.75 - Omicron
Origin:	SARS Coronavirus-2 (SARS-CoV-2), SARS CoV-2 Omicron
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SARS-CoV-2 Spike S1 protein is labelled with His tag.
Drodust Dataila	

Product Details

Purpose:	SARS-CoV-2 Spike S1 Protein, His Tag (BA.2.75/Omicron)		
Sequence:	Val 16 - Arg 685		
Characteristics:	SARS-CoV-2 Spike S1, His Tag (BA.2.75/Omicron) is expressed from human 293 cells		
	(HEK293). It contains AA Val 16 - Arg 685 (Accession # QHD43416.1 (T19I, LPP24-26del, A27S,		
	G142D, K147E, W152R, F157L, I210V, V213G, G257S, G339H, S371F, S373P, S375F, T376A,		
	D405N, R408S, K417N, N440K, G446S, N460K, S477N, T478K, E484A, Q498R, N501Y, Y505H,		
	D614G, H655Y, N679K, P681H). The spike mutations are identified on the SARS-CoV-2 Omicror		
	variant (Pango lineage: BA.2.75).		
Purity:	95,00 %		
Endotoxin Level:	1.0 EU per μg		

Target Details

Target:	SARS-CoV-2 Spike S1	
Abstract:	SARS-CoV-2 Spike S1 Products	
Target Type:	Viral Protein	
Background:	Synonyms:Spike,S1 protein,Spike glycoprotein Subunit1,S glycoprotein Subunit1,Spike protein	
	S1,Description:It's been reported that Coronavirus can infect the human respiratory epithelial	
	cells through interaction with the human ACE2 receptor. The spike protein is a large type I	
	transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor	
	binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2	
	contains basic elements needed for the membrane fusion. The S protein plays key parts in the	
	induction of neutralizing-antibody and T-cell responses, as well as protective immunity.	
Molecular Weight:	76.8 kDa	
Application Details		
Comment:	This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of	
	76.8 kDa. The protein migrates as 110-130 kDa under reducing (R) condition (SDS-PAGE) due to	
	glycosylation.	
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Buffer:	PBS, 0.2M Arginine, pH 7.4	
Storage:	-20 °C	
Storage Comment:	-20°C	