



[Go to Product page](#)

Datasheet for ABIN7316979

## SARS-CoV-2 Spike Protein (BA.5 - Omicron, Trimer) (His tag)

### Overview

Quantity:	200 µg
Target:	SARS-CoV-2 Spike
Protein Characteristics:	BA.5 - Omicron, Trimer
Origin:	SARS Coronavirus-2 (SARS-CoV-2), SARS CoV-2 Omicron
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This SARS-CoV-2 Spike protein is labelled with His tag.

### Product Details

Purpose:	SARS-CoV-2 Spike Trimer Protein, His Tag (BA.5/Omicron) (Trehalose free) (MALS verified)
Sequence:	Val 16 - Pro 1213
Characteristics:	SARS-CoV-2 Spike Trimer, His Tag (BA.5/Omicron) is expressed from human 293 cells (HEK293). The spike mutations are identified on the SARS-CoV-2 Omicron variant (Pango lineage: BA.5, GISAID clade: GRA). The recombinant protein is expressed from human 293 cells (HEK293) with T4 fibrin trimerization motif and a polyhistidine tag at the C-terminus. Proline substitutions (F817P, A892P, A899P, A942P, K986P, V987P) and alanine substitutions (R683A and R685A) are introduced to stabilize the trimeric prefusion state of SARS-CoV-2 S protein and abolish the furin cleavage site, respectively.
Purity:	95,00 %
Endotoxin Level:	1.0 EU per µg
Grade:	MALS verified

## Target Details

---

Target: SARS-CoV-2 Spike

---

Abstract: [SARS-CoV-2 Spike Products](#)

---

Background: Synonyms:Spike,S protein,Spike glycoprotein,S glycoprotein,Description:It's been reported that SARS-CoV-2 can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion.The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

---

Molecular Weight: 137.8 kDa

---

## Application Details

---

Comment: This protein carries a polyhistidine tag at the C-terminus. The protein has a calculated MW of 137.8 kDa. The protein migrates as 160-190 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

---

Restrictions: For Research Use only

---

## Handling

---

Format: Liquid

---

Buffer: PBS

---

Storage: -80 °C

---

Storage Comment: -70°C

---