

## Datasheet for ABIN7317123 **CCL8 Protein (His tag)**



[Go to Product page](#)

### Overview

Quantity:	50 µg
Target:	CCL8
Origin:	Human
Source:	Yeast
Protein Type:	Recombinant
Purification tag / Conjugate:	This CCL8 protein is labelled with His tag.

### Product Details

Purpose:	Recombinant Human CCL8/MCP-2 Protein (Yeast, His Tag)
Sequence:	Gln24-Pro99
Characteristics:	A DNA sequence encoding the human CCL8 (NP_005614.2) (Gln24-Pro99) was expressed with a polyhistidine tag at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.

### Target Details

Target:	CCL8
Alternative Name:	CCL8/MCP-2 ( <a href="#">CCL8 Products</a> )
Background:	Background: Chemokines are a family of small chemotactic cytokines, or proteins secreted by cells. Chemokines share the same structure similarities such as small size, and the presence of four cysteine residues in conserved locations in order to form their 3-dimensional shape. Some of the chemokines are considered pro-inflammatory which can be induced to recruit cells of the

## Target Details

immune system to a site of infection during an immune response, while others are considered homeostatic and are implied in controlling the migration of cells during normal processes of tissue maintenance and development. There are four members of the chemokine family: C-C kemokines, C kemokines, CXC kemokines and CX3C kemokines. The C-C kemokines have two cysteines nearby the amino terminus. There have been at least 27 distinct members of this subgroup reported for mammals, called C-C chemokine ligands-1 to 28. Chemokine ligand 8 (CCL8), also known as monocyte chemoattractant protein 2 (MCP-2), is a small cytokine belonging to the C-C chemokine family. CCL8 functions to activate different immune cells, including mast cells, eosinophils and basophils which are involved in allergic responses, monocytes, and T cells and NK cells which are involved in the inflammatory response. CCL8's ability achieves by binding to different cell surface receptors termed chemokine receptors including CCR1, CCR2B and CCR5. It has been reported that CCL8 is a potent inhibitor of HIV-1 by virtue of its binding to CCR5 which is one of the major co-receptors for HIV-1.

Synonym: C-C Motif Chemokine 8, HC14, Monocyte Chemoattractant Protein 2, Monocyte Chemotactic Protein 2, MCP-2, Small-Inducible Cytokine A8, CCL8, MCP2, SCYA10, SCYA8,HC14

Molecular Weight: 10.3 kDa

NCBI Accession: [NP\\_005614](#)

## Application Details

Restrictions: For Research Use only

## Handling

Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.