



[Go to Product page](#)

Datasheet for ABIN7317521  
**CD200R1L Protein (His tag)**

Overview

Quantity:	100 µg
Target:	CD200R1L
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD200R1L protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human CD200R1L/CD200R2/CD200RLa Protein (His Tag)
Sequence:	Met 1-Leu239
Characteristics:	A DNA sequence encoding the human CD200R1L (AAT00538.1) extracellular domain (Met1-Leu239) was expressed, fused with a polyhistidine tag at the C-terminus.
Purity:	> 95 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.

Target Details

Target:	CD200R1L
Alternative Name:	CD200R1L/CD200R2/CD200RLa ( <a href="#">CD200R1L Products</a> )
Background:	Background: Cell surface glycoprotein CD200 receptor 2, also known as Cell surface glycoprotein CD200 receptor 1-like, Cell surface glycoprotein OX2 receptor 2, CD200 receptor-like 2, CD200Rla, CD200R1L and CD200R2, is a single-pass type I membrane protein which

## Target Details

---

belongs to the CD200R family. CD200R1L / CD200R2. It contains one Ig-like C2-type (immunoglobulin-like) domain and one Ig-like V-type (immunoglobulin-like) domain. CD200 is a transmembrane protein delivering immunoregulatory signals after engagement of CD200R. A family of CD200Rs exist ( CD200R1, CD200R2, CD200R3, CD200R4 ) with different tissue expression and functional activity. In the presence of anti-CD200R2 / CD200R3 monoclonal antibodies (mAbs), bone-marrow cells cultured in the presence of (interleukin [IL]-4+granulocyte-macrophage colony-stimulating factor) differentiate into dendritic cells (DCs), which induce CD4+CD25+ Treg. Interaction between the relatively ubiquitously expressed molecule CD200 and one of its receptors, CD200R1, resulted in direct suppression of alloreactivity, engagement of alternate receptors led instead to altered differentiation of dendritic cells (DCs) from marrow precursors, which could in turn foster development of Foxp3(+) regulatory T cells. Unlike anti-CD200R1, anti-CD200R2 both promotes development of DCs with capacity to induce Treg and directly augments thymocyte production of Treg.

Synonym: CD200R2;CD200RLa

---

Molecular Weight: 25.2 kDa

## Application Details

---

Restrictions: For Research Use only

## Handling

---

Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.