

Datasheet for ABIN7317525 **PVRL1 Protein (Fc Tag)**



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	PVRL1
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This PVRL1 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Human CD111/Nectin-1 Protein (Fc Tag)(Active)
Sequence:	Met 1-Thr 334
Characteristics:	A DNA sequence encoding the human PVRL1 isoform 1 (NP_002846.3) extracellular domain (Met 1-Thr 334) was fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 98 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized rh PVRL3 at 1 µg/ml (100 µl/well) can bind recombinant human Nectin-1 with a linear range of 0.156-5 ng/ml.

Target Details

Target:	PVRL1
---------	-------

Target Details

Alternative Name: CD111/Nectin-1 ([PVRL1 Products](#))

Background: Poliovirus receptor-related 1 (herpesvirus entry mediator C, nectin-1, CD111), also known as PVRL1 is a cell adhesion molecule belonging to the immunoglobulin superfamily that can bind to virion glycoprotein D (gD) to mediate entry of herpes simplex viruses (HSV) and pseudorabies virus (PRV). CD111/Nectin-1/PVRL1 colocalizes with E-cadherin at adherens junctions in epithelial cells. The disruption of cell junctions can result in the redistribution of nectin-1. To determine whether disruption of junctions by calcium depletion influenced the susceptibility of epithelial cells to viral entry, Madin-Darby canine kidney cells expressing endogenous nectin-1 or transfected human nectin-1 were tested for the ability to bind soluble forms of viral gD and to be infected by HSV and PRV, before and after calcium depletion. It has been revealed that binding of HSV and PRV gD was localized to adherens junctions in cells maintained in normal medium but was distributed, along with nectin-1, over the entire cell surface after calcium depletion. Both the binding of gD and the fraction of cells that could be infected by HSV-1 and PRV were enhanced by calcium depletion. Taken together, CD111/Nectin-1/PVRL1 confined to adherens junctions in epithelial cells is not very accessible to virus, whereas dissociation of cell junctions releases nectin-1 to serve more efficiently as an entry receptor.

Synonym: Poliovirus Receptor-Related Protein 1, Herpes Virus Entry Mediator C, Herpesvirus Entry Mediator C, HveC, Herpesvirus Ig-Like Receptor, HIgR, Nectin-1, CD111, PVRL1, HVEC, PRR1,ED4,HIgR,HV1S,HVEC,nectin-1,OFC7,PRR,PVRR,PVRR1,SK-12

Molecular Weight: 61 kDa

NCBI Accession: [NP_002846](#)

Pathways: [Cell-Cell Junction Organization](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C,-20 °C,-80 °C

Handling

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.