

# Datasheet for ABIN7317532

# **SNAP25 Protein (His tag)**



_						
	V	$\triangle$	r۱	/1	$\triangle$	Λ/
	' V '		ΙV			v v

Quantity:	100 μg
Target:	SNAP25
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This SNAP25 protein is labelled with His tag.

### **Product Details**

Purpose:	Recombinant Human SNAP25/SUP Protein (His Tag)	
Sequence:	Met 1-Gly 206	
Characteristics:	A DNA sequence encoding the human SNAP25 (P60880-1) (Met 1-Gly 206) was expressed, with a polyhistide tag at the N-terminus.	
Purity:	> 90 % as determined by reducing SDS-PAGE.	

## **Target Details**

Target:	SNAP25	
Alternative Name:	SNAP25/SUP (SNAP25 Products)	
Background:	Background: Synaptosomal-associated protein 25, also known as Super protein, Synaptosomal-associated 25 kDa protein, SNAP25 and SNAP, is a cytoplasm and cell membrane protein	
	which belongs to the SNAP-25 family. SNAP25 / SUP contains 2 t-SNARE coiled-coil homology	
	domains. SNAP25 / SUP is a membrane bound protein anchored to the cytosolic face of	

membranes via palmitoyl side chains in the middle of the molecule. SNAP25 / SUP protein is a component of the SNARE complex, which is proposed to account for the specificity of membrane fusion and to directly execute fusion by forming a tight complex that brings the synaptic vesicle and plasma membranes together. SNAP25 / SUP is a Q-SNARE protein contributing two  $\alpha$ -helices in the formation of the exocytotic fusion complex in neurons where it assembles with syntaxin-1 and synaptobrevin. SNAP25 / SUP is involved in the molecular regulation of neurotransmitter release. It may play an important role in the synaptic function of specific neuronal systems. SNAP25 / SUP associates with proteins involved in vesicle docking and membrane fusion. SNAP25 / SUP regulates plasma membrane recycling through its interaction with CENPF. SNAP25 / SUP inhibits P/Q- and L-type voltage-gated calcium channels located presynaptically and interacts with the synaptotagmin C2B domain in Ca2+-independent fashion. In glutamatergic synapses SNAP25 / SUP decreases the Ca2+ responsiveness, while it is naturally absent in GABAergic synapses.

Synonym: bA416N4.2;dJ1068F16.2;RIC-4;RIC4;SEC9;SNAP;SNAP-25

Molecular Weight:

24.8 kDa

Pathways:

Positive Regulation of Peptide Hormone Secretion, Hormone Transport, Synaptic Vesicle Exocytosis, Dicarboxylic Acid Transport

#### **Application Details**

Restrictions:

For Research Use only

### Handling

Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from sterile PBS, pH 7.4	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted	
	samples are stable at < -20°C for 3 months.	