.-online.com antibodies

Datasheet for ABIN7317543 NKp44/NCR2 Protein (His tag)



Overview	
Quantity:	100 µg
Target:	NKp44/NCR2 (NCR2)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This NKp44/NCR2 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human NCR2/NKp44/CD336 Protein (His Tag)
Sequence:	Met 1-Pro 190
Characteristics:	A DNA sequence encoding the human NCR2 (NP_004819.2) extracellular domain (Met 1-Pro 190) was fused with a polyhistidine tag at the C-terminus.
Purity:	> 88 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per μ g as determined by the LAL method.

Target Details

Target:	NKp44/NCR2 (NCR2)
Alternative Name:	NCR2/NKp44/CD336 (NCR2 Products)
Background:	Background: Natural cytotoxicity triggering receptor 2 (NCR2), also known as Natural killer cell p44-related protein (NKp44), or CD336, is a member of the natural cytotoxicity receptor (NCR)
	family, which composed of one Ig-like extracellular domain, a transmembrane segment, and a

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7317543 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	cytoplasmic domain. It is a novel transmembrane glycoprotein belonging to the
	Immunoglobulin superfamily characterized by a single extracellular V-type domain. The
	cytoplasmic domain of NKp44 also contains a sequence that matches the immunoreceptor
	tyrosine-based inhibitory motif (ITIM) consensus. This Cytotoxicity-activating receptor that may
	contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell
	lysis. NKp44 is selectively expressed by IL-2-activated NK cells and may contribute to the
	increased efficiency of activated NK cells to mediate tumor cell lysis. Tumor cell recognition of
	the mutated NKp44 proteins was significantly reduced and correlated with their lower
	recognition of heparin.
	Synonym: CD336,dJ149M18.1,LY95,NK-p44,NKP44,RP1-149M18.2
Molecular Weight:	20 kDa
NCBI Accession:	NP_004819
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.