

Datasheet for ABIN7317575

Hsc70 Protein (His tag)



Overview

Quantity:	100 μg
Target:	Hsc70 (HSPA8)
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This Hsc70 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human HSPA8/HSC70 Protein (His Tag)
Sequence:	Met 1-Asp 646
Characteristics:	A DNA sequence encoding the human HSPA8 (P11142-1) (Met 1-Asp 646) was expressed, with a polyhistide tag at the N-terminus.
Purity:	> 90 % as determined by reducing SDS-PAGE.

Target Details

Target:	Hsc70 (HSPA8)
Alternative Name:	HSPA8/HSC70 (HSPA8 Products)
Background:	Background: HSPA8, also known as HSC70, is a member of the heat shock protein family due to homology with other heat shock proteins. The heat shock protein 70 family is comprised by
	both heat-inducible and constitutively expressed members. The latter are called heat-shock
	cognate proteins. HSPA8 belongs to the heat-shock cognate subgroup. Members of the human

heat-shock protein multigene family have several highly conserved proteins with structural and functional properties in common, but vary in the extent of their inducibility in response to metabolic stress. HSPA8 is constitutively expressed and performs functions related to normal cellular processes. This protein binds to nascent polypeptides to facilitate correct protein folding. It also functions as an ATPase in the disassembly of clathrin-coated vesicles during transport of membrane components through the cell. Two alternatively spliced variants have been characterized to date. HSPA8 acts as a repressor of transcriptional activation. It inhibits the transcriptional coactivator activity of CITED1 on Smad-mediated transcription. Isoform 2 may function as an endogenous inhibitory regulator of HSC70 by competing the cochaperones. It also is a ATPase that works with auxilin to remove clathrin coated vesicles. In neurons, synaptojanin is also an important protein involved in vesicle uncoating.

Synonym: HEL-33,HEL-S-72p,HSC54,HSC70,HSC71,HSP71,HSP73,HSPA10,LAP-1,LAP1,NIP71

Molecular Weight:

72.4 kDa

Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, 10 % glycerol, pH 7.5
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.