

Datasheet for ABIN7317581

**EPH Receptor A4 Protein (EPHA4) (His tag,Fc Tag)**[Go to Product page](#)

## Overview

Quantity:	100 µg
Target:	EPH Receptor A4 (EPHA4)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This EPH Receptor A4 protein is labelled with His tag,Fc Tag.

## Product Details

Purpose:	Recombinant Human EphA4 Protein (His & Fc Tag)(Active)
Sequence:	Met 1-Thr 547
Characteristics:	A DNA sequence encoding the human EPHA4 (NP_004429.1) extracellular domain (Met 1-Thr 547) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Purity:	> 92 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized human EPHA5 at 20 µg/ml (100 µl/well) can bind human EFNA4-Fc with a linear ranger of 1.28-32 ng/ml.

## Target Details

Target:	EPH Receptor A4 (EPHA4)
---------	-------------------------

## Target Details

---

Alternative Name: EphA4 ([EPHA4 Products](#))

---

Background: Eph receptor A4 (ephrin type-A receptor 4), also known as EphA4, belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family which 16 known receptors (14 found in mammals) are involved: EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHA9, EPHA10, EPHB1, EPHB2, EPHB3, EPHB4, EPHB5, EPHB6. The Eph family of receptor tyrosine kinases (comprising EphA and EphB receptors) has been implicated in synapse formation and the regulation of synaptic function and plasticity<sup>6</sup>. EphA4 is enriched on dendritic spines of pyramidal neurons in the adult mouse hippocampus, and ephrin-A3 is localized on astrocytic processes that envelop spines. Eph receptor-mediated signaling, which is triggered by ephrins<sup>7</sup>, probably modifies the properties of synapses during synaptic activation and remodeling. Ephrin receptors are components of cell signalling pathways involved in animal growth and development, forming the largest sub-family of receptor tyrosine kinases (RTKs). The extracellular domain of an EphA4 interacts with ephrin ligands, which may be tethered to neighbouring cells. Ligand-mediated activation of Ephs induce various important downstream effects and Eph receptors have been studied for their potential roles in the development of cancer.

Synonym: Ephrin type-A receptor 4;HEK8; SEK; TYRO1;EPHA4;Tyrosine-protein kinase receptor SEK;Tyrosine-protein kinase TYRO1;EK8;hEK8;EPH-like kinase 8

---

Molecular Weight: 86.5 kDa

---

NCBI Accession: [NP\\_004429](#)

---

Pathways: [RTK Signaling](#)

---

## Application Details

---

Restrictions: For Research Use only

---

## Handling

---

Format: Lyophilized

---

Reconstitution: Please refer to the printed manual for detailed information.

---

Buffer: Lyophilized from sterile PBS, pH 7.4

---

Storage: 4 °C,-20 °C,-80 °C

---

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted

samples are stable at  $< -20^{\circ}\text{C}$  for 3 months.