



[Go to Product page](#)

Datasheet for ABIN7317596 CD97 Protein (CD97) (Fc Tag)

Overview

Quantity:	100 µg
Target:	CD97
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD97 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Human CD97 Protein (Fc Tag)(Active)
Sequence:	Met 1-Gln 398
Characteristics:	A DNA sequence encoding the first 398 amino acids (Met 1-Gln 398) of human CD97 isoform 2 (NP_001775.2) extracellular domain was fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA. Immobilized human CD55 at 2 µg/ml (100 µl/well) can bind human CD97 with a linear ranger of 1.28-32 ng/ml.

Target Details

Target:	CD97
---------	------

Target Details

Alternative Name: CD97 ([CD97 Products](#))

Background: Background: The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. The CD97 is a receptor predominantly expressed in leukocytes and belongs to a new group of seven-span transmembrane molecules, which is also designed EGF-TM7 family. The family members are characterized by an extended extracellular region with several N-terminal epidermal growth factor-like domains two of which contain a calcium binding site. Mature CD 97 has two noncovalently associated subunits and is composed of a large extracellular protein (CD97 alpha) and a seven-membrane spanning protein (CD97 beta). CD97 is considered as a defining feature of G protein-coupled receptors. The effects that lymphocytes and erythrocytes adhere to CD97-transfected COS cells suggest that CD97 has the ability to bind cellular ligands. CD97 alpha has three alternatively spliced isoforms that are related to the calcium binding EGF-like repeats in the microfibril protein fibrillin. Leukocytes strongly positive for CD97 are concentrated at sites of inflammation relative to CD97 expression in normal lymphoid tissues.

Synonym: CD97,TM7LN1

Molecular Weight: 68.2 kDa

NCBI Accession: [NP_001775](#)

Application Details

Restrictions: For Research Use only

Handling

Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.