.-online.com antibodies

Datasheet for ABIN7317620 PRDM2 Protein (GST tag)



Overview

Quantity:	100 µg
Target:	PRDM2
Origin:	Human
Source:	Escherichia coli (E. coli)
Protein Type:	Recombinant
Purification tag / Conjugate:	This PRDM2 protein is labelled with GST tag.

Product Details

Purpose:	Recombinant Human PRDM2/RIZ1 Protein (GST Tag)
Sequence:	Met 1-Ala 200
Characteristics:	A DNA sequence encoding the N-terminal segment of human PRDM2 (NP_036363.2) (Met 1- Ala 200), containing the SET domian, was fused with the GST tag at the N-terminus.
Purity:	> 86 % as determined by reducing SDS-PAGE.

Target Details

Target:	PRDM2
Alternative Name:	PRDM2/RIZ1 (PRDM2 Products)
Background:	Background: PR domain containing 2, with ZNF domain (PRDM2), also known as zinc finger protein RIZ, is a member of histone methyltransferase (HMT) class enzymes that methylate
	lysine residues of histones or proteins. HMTs contain a conserved catalytic core termed the
	SET domain, which shares sequence homology with an independently described sequence

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN7317620 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	motif, the PR domain. PRDM2 contains 8 C2H2-type zinc fingers and a distinct SET domain,
	and is highly expressed in retinoblastoma cell lines and in brain tumors, as well as in a number
	of other cell lines and in brain, heart, skeletal muscle, liver and spleen. PRDM2 is a S-adenosyl-L-
	methionine-dependent histone methyltransferase that specifically methylates 'Lys-9' of histone
	H3, and is identified as a tumor suppressor. It is reported that intact $PR(SET)$ sequence is
	required for tumor suppression functions, mutations in the PR domain caused activity reduction
	in human cancers. Also, S-adenosylhomocysteine or methyl donor deficiency inhibits RIZ1 and
	other H3 lysine 9 methylation activities. PRDM2 may also function as a DNA-binding
	transcription factor. It Binds to the macrophage-specific TPA-responsive element (MTE) of the
	HMOX1 (heme oxygenase 1) gene and act as a transcriptional activator. In addition, PRDM2
	(RIZ) is able to binds to the retinoblastoma protein (RB) and also Interacts with GATA3.
	Synonym: HUMHOXY1,KMT8,MTB-ZF,RIZ,RIZ1,RIZ2
Molecular Weight:	49.6 kDa
NCBI Accession:	NP_036363
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile 20 mM Tris, 150 mM NaCl, 0.5 mM DTT, 0.5 mM GSH, pH 8.0
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.