

## Datasheet for ABIN7317760

# **FLT4 Protein (His tag)**



#### Overview

Quantity:	100 μg	
Target:	FLT4	
Origin:	Human	
Source:	HEK-293 Cells	
Protein Type:	Recombinant	
Biological Activity:	Active	
Purification tag / Conjugate:	This FLT4 protein is labelled with His tag.	

#### **Product Details**

Purpose:	Recombinant Human VEGFR3/FLT4 Protein (His Tag)(Active)
Sequence:	Met 1-Ile 776
Characteristics:	A DNA sequence encoding the extracellular domain (Met 1-Ile 776) of human VEGFR3 (NP_002011.2) was expressed with a C-terminal polyhistidine tag.
Purity:	> 97 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its binding ability in a functional ELISA.Immobilized human VEGF-C at 10 $\mu$ g/mL (100 $\mu$ l/well) can bind human VEGFR3-his. The EC50 of human VEGFR3-his is 0.011 $\mu$ g/mL. Measured by its ability to bind human VEGF-D in a functional ELISA.

## **Target Details**

Target:	FLT4	

# **Target Details**

Alternative Name:	VEGFR3/FLT4 (FLT4 Products)	
Background:	Background: Vascular endothelial growth factor receptor 3 (VEGFR3), also known as FLT-4, together with the other two members VEGFR1 (FLT-1) and VEGFR2 (KDR/Flk-1) are receptors for vascular endothelial growth factors (VEGF) and belong to the class III subfamily of receptor tyrosine kinases (RTKs). The VEGFR3 protein is expressed mainly on lymphatic vessels but it is also up-regulated in tumor angiogenesis. Mutations in VEGFR3 have been identified in patients with primary lymphoedema. The VEGF-C/VEGF-D/VEGFR3 signaling pathway may provide a target for antilymphangiogenic therapy in prostate cancer, breast cancer, gastric cancer, lung cancer, non-small cell lung cancer (NSCLC), and so on.Immune  Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy  Synonym: FLT-4;FLT-41;FLT41;LMPH1A;PCL;VEGF Receptor 3;VEGFR-3;VEGFR3	
Molecular Weight:	86 kDa	
NCBI Accession:	NP_002011	
Pathways:	RTK Signaling, Signaling Events mediated by VEGFR1 and VEGFR2, VEGF Signaling	
Application Details		
Restrictions:	For Research Use only	
Handling		
Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from sterile PBS, pH 7.4	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	