

Datasheet for ABIN7317812

LIFR Protein (His tag)

[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	LIFR
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This LIFR protein is labelled with His tag.

Product Details

Purpose:	Recombinant Human LIFR/CD118 Protein (His Tag)(Active)
Sequence:	Met 1-Ser 833
Characteristics:	A DNA sequence encoding the extracellular domain of human LIFR (NP_001121143.1) (Met 1-Ser 833) with a C-terminal polyhistidine tag was expressed.
Purity:	> 90 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by its ability to bind human LIF-Fc in a functional ELISA.

Target Details

Target:	LIFR
Alternative Name:	LIFR/CD118 (LIFR Products)

Target Details

Background:	<p>Background: LIFR (leukemia inhibitory factor receptor) belongs to the family of cytokine receptors. LIFR forms a high-affinity receptor complex with gp130, which mediates the activity of LIF (leukemia inhibitory factor) and thus affects the differentiation, proliferation, and survival of a wide variety of cells in the adult and the embryo. Besides LIF, LIFR can also bind to and activate CNTF (ciliary neurotrophic factor) and CLC (cardiotrophin like cytokine). Evidence showed that in the retina, LIFR activating LIF, CT-1 and cardiotrophin like cytokine (CLC) are strongly upregulated in response to preconditioning with bright cyclic light leading to robust activation of signal transducer and activator of transcription-3 (STAT3) in a time-dependent manner. Further, blocking LIFR activation during preconditioning using a LIFR antagonist (LIF05) attenuated the induced STAT3 activation and also resulted in reduced preconditioning-induced protection of the retinal photoreceptors. These data demonstrate that LIFR and its ligands play an essential role in endogenous neuroprotective mechanisms triggered by preconditioning-induced stress. LIFR was newly found to be a suppressor of hepatocellular carcinoma (HCC), one of the world's top five causes of cancer-related deaths.</p> <p>Synonym: CD118,LIF-R,SJS2,STWS,SWS</p>
-------------	--

Molecular Weight:	91 kDa
-------------------	--------

NCBI Accession:	NP_001121143
-----------------	------------------------------

Pathways:	JAK-STAT Signaling , Growth Factor Binding
-----------	--

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
---------	-------------

Reconstitution:	Please refer to the printed manual for detailed information.
-----------------	--

Buffer:	Lyophilized from sterile PBS, pH 7.4
---------	--------------------------------------

Storage:	4 °C,-20 °C,-80 °C
----------	--------------------

Storage Comment:	<p>Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.</p> <p>Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>
------------------	--