



[Go to Product page](#)

Datasheet for ABIN7317868
Cadherin 5 Protein (CDH5) (His tag,Fc Tag)

Overview

Quantity:	100 µg
Target:	Cadherin 5 (CDH5)
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This Cadherin 5 protein is labelled with His tag,Fc Tag.

Product Details

Purpose:	Recombinant Human VE-Cadherin/CDH5 Protein (His & Fc Tag)(Active)
Sequence:	Met 1-Gln 593
Characteristics:	A DNA sequence encoding the extracellular domain of human CDH5 (NP_001786.2) (Met 1-Gln 593) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.
Purity:	> 85 % as determined by reducing SDS-PAGE.
Endotoxin Level:	< 1.0 EU per µg as determined by the LAL method.
Biological Activity Comment:	Measured by the ability of the immobilized protein to support the adhesion of MCF-7 human breast adenocarcinoma cells. When 5 x 10 ⁴ cells/well are added to Recombinant Human Cadherin-5 coated plates (0.8 µg/mL with 100 µL/well), approximately >50% will adhere after 1 hour at 37°C.

Target Details

Target:	Cadherin 5 (CDH5)
Alternative Name:	VE-Cadherin/CDH5 (CDH5 Products)
Background:	<p>Background: Cadherins (Calcium dependent adhesion molecules) are a class of transmembrane proteins. Cadherin-5, also known as VE-cadherin, CDH5 and CD144, an endothelial specific cell-cell adhesion molecule, plays a pivotal role in the formation, maturation and remodeling of the vascular wall. VE-Cadherin is widely considered to be specific for vascular endothelia in which it is either the sole or the predominant cadherin, often co-existing with N-cadherin. This specificity of VE-cadherin for vascular endothelial cells is important not only in blood and lymph vessel biology and medicine, but also for cell-type-based diagnoses, notably those of metastatic tumors. As a classical cadherin, VE-Cadherin links endothelial cells together by homophilic interactions mediated by its extracellular part and associates intracellularly with the actin cytoskeleton via catenins. Mechanisms that regulate VE-cadherin-mediated adhesion are important for the control of vascular permeability and leukocyte extravasation. In addition to its adhesive functions, VE-Cadherin regulates various cellular processes such as cell proliferation and apoptosis and modulates vascular endothelial growth factor receptor functions. Consequently, VE-cadherin is essential during embryonic angiogenesis.</p> <p>Synonym: 7B4,CD144</p>
Molecular Weight:	92 kDa
NCBI Accession:	NP_001786
Pathways:	Cell-Cell Junction Organization, Signaling Events mediated by VEGFR1 and VEGFR2

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted

samples are stable at < -20°C for 3 months.